

MUCOSOLVAN cough syrup 30 mg/5 ml contains propylene glycol.
This medicine contains 28.75 mg propylene glycol in 5 ml solution. If your baby is under 4 weeks of age ask your doctor or pharmacist before giving this medicine to your baby; in particular if your baby is given other medicines that contain propylene glycol or alcohol.

3. How to take MUCOSOLVAN cough syrup 30 mg/5 ml

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. The recommended dose is:

Children up to 2 years of age: on medical advice only
1.25 ml of solution should be taken twice daily (equivalent to 15 mg ambroxol hydrochloride per day)

Children from 2 to 5 years of age:
1.25 ml of solution should be taken three times daily (equivalent to 22.5 mg ambroxol hydrochloride per day)

Children from 6 to 12 years of age:
2.5 ml of solution should be taken two to three times daily (equivalent to 30 - 45 mg ambroxol hydrochloride per day)

Adults and adolescents from 12 years of age:
The usual dosage is 5 ml of solution three times daily (equivalent to 90 mg ambroxol hydrochloride per day) for the first 2 - 3 days and 5 ml of solution twice daily (equivalent to 60 mg ambroxol hydrochloride per day) thereafter.

For adults and adolescents from 12 years of age, the therapeutic effect may be enhanced by increasing the dosage to 10 ml twice daily (equivalent to 120 mg ambroxol hydrochloride per day).

Dosage scheme	
Age	Daily dose
0 - 2 years:	
On medical advice only	2 x 1.25 ml
2 - 5 years	3 x 1.25 ml
6 - 12 years	2 - 3 x 2.5 ml
From 12 years:	
On the first 2 - 3 days	3 x 5 ml
Thereafter	2 x 5 ml

Method of administration and duration of treatment
MUCOSOLVAN cough syrup 30 mg/5 ml is an oral solution. MUCOSOLVAN cough syrup 30 mg/5 ml can be taken with or without food and is dispensed with the aid of the measuring cup provided. Please clean the measuring cup after each use by rinsing with warm water. You must talk to a doctor if you feel worse or if you do not feel better after 4 - 5 days. If you feel that the effect of MUCOSOLVAN cough syrup 30 mg/5 ml is too strong or too weak, talk to your doctor or pharmacist.

If you take more MUCOSOLVAN cough syrup 30 mg/5 ml than you should
No specific symptoms of overdose have been reported to date. The symptoms observed in cases of accidental overdose or medication error are consistent with the side effects which may occur at recommended doses (see section 4). In case of overdose, please consult a doctor as symptomatic treatment may be required.

If you forget to take MUCOSOLVAN cough syrup 30 mg/5 ml
If you forget to take MUCOSOLVAN cough syrup 30 mg/5 ml

or take too small a dose, wait until it is time to take your next dose and then continue taking MUCOSOLVAN cough syrup 30 mg/5 ml as described in the dosage instructions above.

If you stop taking MUCOSOLVAN cough syrup 30 mg/5 ml
If you stop treatment early, your symptoms may get worse.

4. Possible side effects

Like all medicines, MUCOSOLVAN cough syrup 30 mg/5 ml can cause side effects, although not everybody gets them. The following categories are used when reporting the frequencies of side effects:

Very common:	May affect more than 1 in 10 treated patients
Common:	May affect up to 1 in 10 treated patients
Uncommon:	May affect up to 1 in 100 treated patients
Rare:	May affect up to 1 in 1,000 treated patients
Very rare:	May affect up to 1 in 10,000 treated patients
Not known:	Frequency cannot be estimated from the available data

Side effects

Immune system disorders

Rare: Hypersensitivity reactions
Not known: Anaphylactic reactions including anaphylactic shock, angioedema (rapidly developing swelling of the skin, subcutaneous, mucosa or submucosal tissues) and pruritus

Skin and subcutaneous tissue disorders

Rare: Rash, urticaria
Not known: Severe cutaneous adverse reactions (including erythema multiforme, Stevens-Johnson syndrome/toxic epidermal necrolysis and acute generalized exanthematous pustulosis).

Nervous system disorders

Common: Taste disturbances

Gastrointestinal disorders

Common: Nausea, feeling of numbness in the mouth
Uncommon: Vomiting, diarrhoea, indigestion, abdominal pain, dry mouth

Rare: Dry throat
Very rare: Increased salivation

Respiratory, thoracic and mediastinal disorders

Common: Feeling of numbness in the throat
Not known: Shortness of breath (as symptom of a hypersensitivity reaction)

General disorders and administration site conditions

Uncommon: Fever, mucous membrane reactions

Action to be taken in the event of side effects

At the first signs of a hypersensitivity reaction, stop taking MUCOSOLVAN cough syrup 30 mg/5 ml.

Reporting of side effects

If you get any side effects, talk to your doctor, your pharmacist or your nurse. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store MUCOSOLVAN cough syrup 30 mg/5 ml

KEEP THIS MEDICINE OUT OF THE SIGHT AND REACH OF CHILDREN

Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Storage conditions

Do not store above 25°C.

Shelf life after first opening the bottle

The shelf life after first opening the bottle is 6 months.

Do not throw away any medicines via wastewater (e.g. not via the toilet or the sink). Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Further information

What MUCOSOLVAN cough syrup 30 mg/5 ml contains:

The active substance is: ambroxol hydrochloride
5 ml of solution contains 30 mg ambroxol hydrochloride.

The other ingredients are: Benzoic acid, hydroxyethylcellulose, sucralose, flavouring agents (containing propylene glycol), purified water.

What MUCOSOLVAN cough syrup 30 mg/5 ml looks like and contents of the pack:

MUCOSOLVAN cough syrup 30 mg/5 ml is a colourless liquid with a fruity odor in an amber glass bottle with child-resistant screw closure.

MUCOSOLVAN cough syrup 30 mg/5 ml is available in packs containing 100 ml or 250 ml.

Not all pack sizes may be marketed in your country.

Marketing Authorization Holder

A. Nattermann & Cie. GmbH
Brüningstrasse 50
65929 Frankfurt am Main, Germany


Manufacturer

Delpharm Reims
10 Rue Colonel Charbonneaux
51100 Reims, France

This leaflet was last revised in: July 2022.

Child-resistant bottle cap:

1. Remove measuring cup from cap
2. To open, push down on cap
3. At the same time, twist cap in the direction of the arrow




تَمَت مراجعة هذه النشرة للمرة الأخيرة في: تمّوز/يوليو 2022.

غطاء القنينة المقاوم للأطفال:

1. إنزع كوب القياس عن الغطاء

2. للفتح، إضغط على الغطاء

3. في الوقت ذاته، أدر الغطاء باتجاه السهم



التدابير المضادة في حالة حصول تأثيرات جانبية
في حالة ظهور العلامات الأولى لارتكاسات فرط الحساسية، يجب التوقف عن تناول ميوكوسولفان شراب للسعال 30 ملغ/5 مل.

الإفادة عن التأثيرات الجانبية
إذا لاحظت أي تأثيرات جانبية، فعليك استشارة طبيبك أو الصيدلي أو الممرض/ة. ينطبق هذا أيضًا على أي تأثيرات جانبية غير مدرجة في هذه النشرة. بالإفادة عن التأثيرات الجانبية، يمكنك المساعدة على تزويد المزيد من المعلومات حول سلامة هذا الدواء.

5. كيف يُحفظ ميوكوسولفان شراب للسعال 30 ملغ/5 مل
إحفظ هذا الدواء بعيدًا عن نظر الأطفال ومتناولهم. لا تستعمل هذا الدواء بعد تاريخ انتهاء الصلاحية المدوّن على اللصاقة وعلبة الكرتون بعد عبارة "EXP". يشير تاريخ انتهاء الصلاحية إلى اليوم الأخير من الشهر المذكور.

شروط الحفظ
يحفظ في درجة حرارة لا تتجاوز 25 درجة مئوية.

مدة الحفظ بعد فتح القنينة
مدة الحفظ بعد فتح القنينة هي 6 أشهر. لا ينبغي رمي أيّ الدواء مع مياه الصرف الصحي (مثلاً ليس عبر المراض أو المعلقة). إسأل الصيدلي عن كيفية التخلص من الأدوية التي لم تعد تستعملها، فمن شأن هذه الإجراءات حماية البيئة.

6. معلومات إضافية
ماذا يحتوي ميوكوسولفان شراب للسعال 30 ملغ/5 مل:
المادة الفاعلة هي: هيدروكلوريد أمبروكسول.

يحتوي 5 مل من المحلول على 30 ملغ من هيدروكلوريد أمبروكسول. المكونات الأخرى هي: حمض البنزويك، هيدروكسي إيثيل سلولوز، سكرالوز، عوامل منكهة (تحتوي بروبيلين غليكول)، ماء منقى.

كيف هو شكل ميوكوسولفان شراب للسعال 30 ملغ/5 مل ومحتويات العلبة
ميوكوسولفان شراب للسعال 30 ملغ/5 مل هو سائل عديم اللون مع رائحة فواكه في قنينة زجاجية لها غطاء يُفتح بشكل لولبي مقاوم للأطفال. ميوكوسولفان شراب للسعال 30 ملغ/5 مل متوفّر في علب تحتوي على 100 مل أو 250 مل. قد لا تكون أحجام العبوات كلها متسوّقة في بلدك.

حامل رخصة التسويق
A. Nattermann & Cie. GmbH
Brüningstrasse 50
65929 Frankfurt am Main, Germany

المصنّع
Delpharm Reims
10 rue Colonel Charbonneaux
51100 Reims, France